

A Possible Relationship between Size and Reproductive Behavior in a Population of *Aplysia punctata* Cuvier, 1803

BY

CARY OTSUKA^{1,3}, YVES ROUGER² AND ETHEL TOBACH¹

INTRODUCTION

AS PART OF A COMPARATIVE STUDY of the behavioral adaptations of *Aplysia* (TOBACH, *et al.*, 1965; LEDERHENDLER, *et al.*, 1975; LEDERHENDLER, *et al.*, 1977; LEDERHENDLER & TOBACH, 1978; TOBACH, 1978), a study of *Aplysia punctata* was carried out in the intertidal waters of Concarneau, France (47°53'22"N, 3°54'10"W). From June 21, 1979, through July 28, 1979, observations were conducted approximately 800m W of the Laboratoire de Biologie Marine, Collège de France. The field site measured approximately 240m² and was enclosed by large rocks covered primarily by *Fucus serratus*. The basin consisted of coarse sand, scattered rocks of different sizes covered with several species of algae. These were, in addition to the *F. serratus*, *Palmaria marina*, *Enteromorpha* spp., *Lomentaria articulata*, *Laminaria digitata*, *Ulva* spp., and *Chondrus* spp. The rocks were usually covered by more than one algal species; the most abundant were *Fucus*, *Enteromorpha* and *Ulva*. The maximum tidal range was approximately 4.3m and during periods of extreme low tides, the entire area became exposed. Surface water temperature ranged from 15° to 20° C with a fairly constant salinity of 34‰.

PROCEDURES

Vegetation, substrate and rocks were systematically searched by sight and touch for animals, for one half hour before and after the lowest point of the

Table 1
Field observations of *Aplysia punctata*.

Date (1979)	Number of animals found ¹	Median		
		Weight (g)	Range	
			From	To
6/21(AM)	18	48	30	75
6/21(PM)	15	35	15	55
6/22	18	38	10	53
6/23	18	42	30	70
6/24	24	40	25	65
6/25	20	45	25	65
6/26	4	45	30	<50
6/27	21	40	<30	65
6/28	14	38	<30	50
6/29	14	48	25	70
6/30	1	55	—	—
7/1	10	45	>25	75
7/4	1	50	—	—
7/7	15	45	35	75
7/8	14	40	>20	70
7/9	10	40	25	75
7/10	12	40	30	65
7/12	23	40	25	50
7/13	30	40	20	65
7/15	10	40	30	65
7/16	3	45	25	45
7/22	5	35	25	50
7/24	12	40	25	65
7/25	4	30	20	40
7/27	1	40	—	—

¹Only animals found for the first time are listed.

¹ Department of Animal Behavior, American Museum of Natural History, New York; and the Biology Department, City College, The City University of New York; for reprints write to E. Tobach

² Institut National pour Recherche Agronomique, Laboratoire de Biologie Marine, Collège de France, Concarneau, Bretagne, France

³ Research supported by grants from the Arts Foundation and the Lerner Fund for Marine Research of the American Museum of Natural History. We wish to thank Director Yves Legal, Collège de France, for his support and cooperation

tide. Once animals were located, records were made of type of contact and weights by volumetric measurements in graduated cylinders to the nearest 5 mL. A Hartner Balance, model LP105, was used to determine the net weights of animals which were also measured volumetrically. The difference in measurements by the two methods was not noteworthy (± 0.5 g) and it was assumed that mL = g. Animals were tagged according to methods developed by Tobach (LEDERHENDLER *et al.*, 1975). Given the restrictions of time and tidal conditions, the data obtained can only be understood within this context. As Table 1 shows, there was a considerable variation in the number of animals found on any day.

RESULTS AND DISCUSSION

Table 1 lists the days on which observations were made, the number of animals tagged and their weights. It should be noted that during the month of August, on 15 days of searching, 2-6 animals were found on only 4 days. Between the 11th and 26th of August (last observation date), no animals were found.

Table 2 shows the number of times animals were found in contact or copulating. In addition, 194 sea hares were found not engaged in either activity.

Table 2

Aplysia punctata copulating or in contact:
Number of times seen.

		Number of animals			
A. Copulating	2	3	4		
	89	7	3 ²		
B. Contact	2	3	4	7	
	22	9	5	2	

²Weights of one chain of 4 not obtained.

Table 3 shows that a significant number of copulating pairs found during the observation period were composed of sperm donors which were smaller than sperm recipients (one-sample Chi-squared test (SIEGEL, 1956), $p < 0.001$). Given the results of this statistical analysis, pairs in which donors were smaller than recipients were classed as "typical," and others as "atypical."

Table 4 shows that the donor in the most frequently found relationship, *i.e.*, "typical" donor (D'), is, indeed, smaller than the recipient in the "typical" relationship (R'). R' is smaller than either of the other animals in the "atypical" pairing, that is, R^a; where "t" stands for "typical" and "a" stands for "atypical." The recipient in the typical pairing (R') is also larger than the recipient in the atypical pairing (R^a).

The lack of a significant difference in weight between R' and D^a would seem to indicate that at a certain weight, these sea hares are as likely to be found acting as sperm donors as sperm recipients.

In the laboratory, 2 sea hares (*Aplysia punctata* Cuvier, 1803), weighing 3 g and 8.7 g were seen copulating, the smaller being the sperm donor. The smaller sea hare was seen acting as sperm donor until it weighed 7.1 g two weeks later. Subsequently both animals assumed both roles and laid eggs. The sea hare that acted as recipient in the early copulations first laid eggs when it weighed 11.6 g. LEDERHENDLER (1978) reported that a pair of *A. dactylomela* first copulated when weighing 58 g and 53 g, the former being the sperm donor. However, subsequent to this first copulation, the smaller of the 2 was the sperm donor on the 3 occasions when copulation took place. The size relationship seems to be similar in the *A. dactylomela*, which appear to attain greater size than *A. punctata* (see CAREFOOT, 1967 for *A. punctata* and Tobach, unpubl. data for Bimini; and LEDERHENDLER *et al.*, 1977, for Puerto Rico for *A. dactylomela*).

The weights of the laboratory *Aplysia punctata* are in a very different range from those observed in the field. However, in the pair described above which was followed daily in the laboratory, the size relationship was consistent with the field data.

These data are preliminary in that the sampling technique was restricted to a particular field population at only one point in their daily activity under one tidal con-

Table 3

Patterns of copulation and individual weights of *Aplysia punctata*.

A. Paired Animals

		Donor smaller than recipient	Donor larger than recipient	Donor equal to recipient
Number of pairs ³		56 ⁴	25 ⁵	8
Weights (g)	Donor:			
	Median	30	50	43
	Range from	20	25	30
	to	70	75	60
	Recipient:			
	Median	50	35	See above
	Range from	30	25	
	to	75	65	

B. Chains of 3 and 4 Animals

Copulatory relationships ⁶		Donor	Donor and recipient	Recipient
Typical	Chain of 3	20	30	40
	of 3	25	30	40
	of 4	30	35	60
			40	
Atypical	Chain of 3	45	40	35
	of 3	60	40	30
Mixed	Chain of 3	40	30	40
	of 3	25	40	35
	of 3	35	35	65
	of 4	30	30	70
			44	

³X² = 37.8; p < 0.001.⁴This relationship, i.e., donor (D) < recipient (R) in weight, is classed as "typical" ("t"), because of the significant X².⁵This relationship, i.e., donor > recipient is classed as "atypical" ("a"), for the same reason.⁶Categories "typical," and "atypical" based on statistical analysis of data for paired animals in "A" above. "Mixed" indicates that donor/recipient was "typical" for one part of the chain and "atypical" in another part.

dition. These findings suggest, however, that the commonly held belief that *Aplysia* assume both roles soon after metamorphosis needs to be re-examined, and that further study is warranted.

Literature Cited

- CAREFOOT, T. H.
1967. Studies on a sublittoral population of *Aplysia punctata*.
Journ. Mar. Biol. Assoc. U. K. 47: 335-350

Table 4

Analysis⁷ of weights of copulating pairs of *Aplysia punctata*.

Copulatory role	Distribution according to median weight (40g) for combined groups			
	$D^t < R^t$ ⁸		$D^a > R^a$ ⁸	
	D^t	R^t	D^a	R^a
Number of animals				
Below median	43	5	3	13
At median	6	7	5	6
Above median	7	44	17	6

⁷Median test (Siegel, 1956).⁸D = donor; R = recipient; t = "typical" relationship; a = "atypical" relationship. $D^t < R^t$; "p" < 0.001 $D^t < D^a$; 0.10 > "p" > 0.05 $D^a < R^a$; "p" < 0.02 D^a and R^a ; do not differ: "p" > 0.10 $R^t > R^a$; "p" < 0.001

LEDERHENDLER, IZJA, LARRY BELL & ETHEL TOBACH

1975. Preliminary observations on the behavior of *Aplysia dactylomela* (Rang, 1828) in Bimini waters. *The Veliger* 17 (4): 347-353; 1 text fig. (1 April 1975)

LEDERHENDLER, IZJA, K. HERRIGES & ETHEL TOBACH

1977. Taxis in *Aplysia dactylomela* (Rang, 1828) to water-borne stimuli from conspecifics. *Anim. Learn. Behav.* 5: 355-358

LEDERHENDLER, IZJA & ETHEL TOBACH

1977. Reproductive roles in the simultaneous hermaphrodite *Aplysia dactylomela*. *Nature* 270: 238-239

SIEGEL, S.

1956. Nonparametric statistics for the behavioral sciences. McGraw-Hill Book Co., New York

TOBACH, ETHEL

1978. Further field notes on the behavior of *Aplysia dactylomela*.*The Veliger* 20 (4): 359-360 (1 April 1978)

TOBACH, ETHEL, PETER GOLD & AMY ZIEGLER

1965. Preliminary observations of the inking behavior of *Aplysia* (*Varria*). *The Veliger* 8 (1): 16-18 (1 July 1965)